

Gene transcription of neuroglobin is upregulated by hypoxia and anoxia in the brain of the anoxia-tolerant turtle *Trachemys scripta*

Sarah L. Milton^{1,*}, Gauri Nayak¹, Peter L. Lutz¹ & Howard M. Prentice²

¹Department of Biological Sciences SC 281, Florida Atlantic University, 777 Glades Road, Boca Raton, FL, 33431, USA; ²Department of Biomedical Sciences, Florida Atlantic University, 777 Glades Road, Boca Raton, FL, 33431, USA

Received 15 February 2006; accepted 21 March 2006
© 2006 National Science Council, Taipei

Key words: anoxia, hypoxia, neuroglobin, *Trachemys scripta*

Summary

Neuroglobin is a heme protein expressed in the vertebrate brain in mammals, fishes, and birds. The physiological role of neuroglobin is not completely understood but possibilities include serving as an intracellular oxygen-carrier or oxygen-sensor, as a terminal oxidase to regenerate NAD⁺ under anaerobic conditions, or involvement in NO or ROS metabolism. As the vertebrate nervous system is particularly sensitive to hypoxia, an intracellular protein that helps sustain cellular respiration would aid hypoxic survival. However, the regulation of Neuroglobin (Ngb) under conditions of varying oxygen is controversial. This study examines the regulation of Ngb in an anoxia-tolerant vertebrate under conditions of hypoxia and anoxia. The freshwater turtle *Trachemys scripta* can withstand complete anoxia for days, and adaptations that permit neuronal survival have been extensively examined. Turtle neuroglobin specific primers were employed in RT-PCR for determining the regulation of neuroglobin mRNA expression in turtles placed in normoxia, hypoxia (4 h), anoxia (1 and 4 h), and anoxia-reoxygenation. Whole brain expression of neuroglobin is strongly upregulated by hypoxia and post-anoxic-reoxygenation in *T. scripta*, with a lesser degree of upregulation at 1 and 4 h anoxia. Our data implicate neuroglobin in mediating brain anoxic survival.

Introduction

Neurons are generally viewed as among the most anoxia sensitive of all cells, though recent studies have shown a wide variation in the capacity of neurons to tolerate hypoxia, reflective of their function and the degree of hypoxia normally encountered. Even the most vulnerable neurons are not defenseless, and the most tolerant are able to withstand extreme periods of complete anoxia and recover fully when oxygen becomes available [1]. The fate of hypoxic or ischemic neurons

depends upon both physiological and molecular events, with pro-survival and apoptotic pathways competing at the transcriptional and post-transcriptional level. These events include the upregulation of various pathways that increase anaerobic metabolism or oxygen delivery, including erythropoietin [2], vascular endothelial growth factor [3] and heat shock proteins [4]. At the physiological level, pro-survival pathways include the opening of ATP-regulated potassium channels [5], and increases in inhibitory neurotransmitters including GABA [6], and adenosine [7, 8].

Not all vertebrate brains, however, are equally sensitive to hypoxia. Freshwater turtles of the genus *Trachemys* are true facultative anaerobes,

*To whom correspondence should be addressed. Fax: +561-297-2749; E-mail: smilton@fau.edu

able to survive from up to 48 h at room temperature to months (during winter hibernation) in the total absence of oxygen [9]. *Trachemys scripta* has been the subject of extensive research into the adaptations that permit neuronal survival without oxygen [1]; the turtle brain is able to decrease its metabolic rate to approximately 10–15% of basal, such that energy utilization is matched to anaerobic energy production. By preventing an energy deficit, the turtle brain avoids the catastrophic drop in ATP levels which, in mammalian neurons, results in the breakdown of cellular ion homeostasis, release of excitatory neurotransmitters, and excitotoxic cellular death [1]. To decrease neuronal energy requirements, *Trachemys* decreases membrane ion permeability (“channel arrest”), inhibits the release of excitatory neurotransmitters such as dopamine [10] and glutamate [11], increases the release of inhibitory compounds including adenosine [12] and GABA [13], and decreases electrical activity [14]. In terms of molecular changes, organ-level alterations of MAP kinases, ERK, and JNK have been reported [15], while work in our lab has shown increased expression of heat shock proteins [16] and the downregulation of Kv channel transcription [17]; alterations in many of these molecular factors have been linked to hypoxic/ischemic survival in the mammalian CNS. Such extended anoxic survival time is not a matter of ectothermy, as other reptiles survive only 20–30 min without oxygen [18] and do not exhibit the same neurological adaptations that permit true anoxic tolerance [19].

As the CNS in most vertebrates, however, is particularly sensitive to O₂ lack, the increased expression of a protein that helps to sustain cellular respiration could increase neuronal survival under the low oxygen conditions of hypoxia or ischemia. Neuroglobin (Ngb), a recently discovered heme protein of the CNS, is a highly conserved protein able to bind oxygen reversibly. First reported in mouse brain [20], Ngb has since been identified as well in fish, amphibians, and birds [21], pointing to its likely presence in all vertebrates. Globins are usually considered either oxygen transporter or O₂ storage proteins, therefore a likely role for Ngb is as a “neuronal myoglobin” transferring oxygen to the mitochondrial respiratory chain in the brain and retina [21]. Other potential roles include acting as a terminal oxidase that regenerates NAD⁺ under anaerobic conditions [22–24], aiding in the detoxification of

reactive oxygen species, or as a sensor to detect cellular oxygen concentrations.

To date, however, it is unclear if Ngb is regulated under varying O₂ levels, with some studies reporting increased expression of Ngb mRNA and protein in cerebral tissue culture after 24 h anoxia-reperfusion [25], while other investigations [26] did not find increased expression in hypoxic (10% O₂) mice *in vivo*. These conflicting results, however, could indicate that Ngb is involved only during acute hypoxia. Increased levels of Ngb have been shown to protect against hypoxic/ischemic injury in both cultured neurons [25] and in an experimental stroke model in rats [27]. Because of the exquisite sensitivity of the mammalian brain to severe hypoxia/anoxia, however, investigation of the regulation of Ngb under such conditions is difficult. The brain of the anoxia-tolerant turtle offers a unique model to identify strategies to enhance the survival of neurons vulnerable to hypoxia/reoxygenation stress, while providing a means to separate protective responses from the mammalian mixture of adaptive and pathological responses [28].

In this study, we examined the *in vivo* expression of Ngb mRNA using RT-PCR in whole *T. scripta* brains in hypoxia (5% O₂), short term (1 h) and long-term (4 h) anoxia, and upon anoxia–reoxygenation.

Material and methods

All experiments were conducted with the approval of Florida Atlantic University Institutional Animal Care and Use Committee. Freshwater turtles (*Trachemys scripta*) weighing 300–500 g obtained from commercial suppliers (Lemberger Reptiles, Oshkosh, WI) were maintained at room temperature (22–23 °C) in freshwater aquaria on a 12 h light/dark cycle. Animals were fed 3× weekly to satiation on commercial turtle food.

Tissue preparation

Five experimental sets of $n=5$ included normoxic controls, anoxic animals exposed to 1 or 4 h anoxia, 4 h anoxia/4 h normoxic recovery, and 4 h hypoxia (5% O₂). For anoxic exposure, animals were individually placed in sealed 2 L plastic chambers at room temperature (23 °C) under

99.99% N₂ (positive pressure flow-through, County Welding, Pompano Beach, FL). Normoxic controls were utilized directly from the aquaria. Hypoxic animals were placed in individual open holding boxes in a hypoxia chamber (Sheldon Manufacturing, Cornelius, OR) under 5% O₂ for 4 h. Chamber PO₂ was determined hourly to ascertain O₂ levels (Cameron Instrument Company, Port Aransas, TX).

Animals were sacrificed by cervical separation and the brains removed into liquid nitrogen in less than 2 min.

RT-PCR

Total RNA was extracted using the TRIzol reagent (Life Technologies, Grand Island, NY) according to the manufacturer's protocol and RNA was subjected to treatment with DNase I to eliminate DNA contamination. Complementary DNA was synthesized from total RNA using primers specific for neuroglobin and actin, respectively. The PCR using *Taq* polymerase comprised denaturation for 7 min, 94 °C, PCR: 40 cycles (Ngb) (1 min, 94 °C; 45 s, 59 °C; 1.0 min, 72 °C) followed by elongation: 10 min, 72 °C or 30 cycles (actin) (1 min 94 °C; 45 s, 55 °C; 1.5 min 72 °C) followed by elongation: 10 min 72 °C.

Primers specific to turtle neuroglobin cDNA were designed from a partial cDNA sequence that was obtained previously by RT-PCR analysis of turtle brain mRNA using degenerate primers homologous to neuroglobin sequences from mouse and zebrafish. The primers employed for PCR were the following: turtle brain specific neuroglobin primers: 5'-GTTGTTTGATCTGGACCCTGAC-3' (forward) and 5'-TTGCCAAGTTGGAGA

GATATT-3' (reverse); actin primers: 5'-CACCAACTGGGACGACATGG-3' (forward) and 5'-GTCGGCCAGCTCGTAGCTCT-3' (reverse) (Table 1).

PCR products were separated by gel electrophoresis, visualized by ethidium bromide and photographed using a digital camera for quantification using National Institute of Health Image J 1.60 software.

For semi-quantitative measurement of neuroglobin transcript levels, RT-PCR signal intensities were calculated as a ratio of levels of PCR products amplified from turtle actin cDNAs. Data are expressed as percent increase above normoxic control.

Statistical analysis: data were analyzed by ANOVA (Dunn's Multiple Comparison test; $p < 0.05$ was considered to be significant).

Results

To establish whether neuroglobin transcription was influenced by oxygen availability, we measured levels of neuroglobin mRNA expression in brain from normoxic turtles and turtles subjected to 4 h hypoxia at 5% O₂, 1 h anoxia, 4 h anoxia, and 4 h anoxia followed by 4 h reoxygenation. Levels of mRNA expression were normalized to actin mRNA control levels.

A clear upregulation of neuroglobin mRNA expression (3.5-fold increase, $p < 0.01$) was seen at 4 h hypoxia whereas a more modest but progressive increase was found in 1 and 4 h anoxia reaching a 2.0-fold increase by 4 h anoxia ($p < 0.05$) (Figures 1 and 2). On reoxygenation following 4 h anoxia high levels of neuroglobin

Table 1. Primer sequences and size of RT-PCR products.

Target gene	Predicted size (bp)	Sense/Antisense	Location, nucleotides
Neuroglobin	193	5'-GTTGTTTGATCTGGACCCTGAC-3'	368–389
		5'-TTGCCAAGTTGGAGAGATATT-3'	559–580
Actin	505	5'-CACCAACTGGGACGACATGG-3'	228–247
		5'-GTCGGCCAGCTCGTAGCTCT-3'	713–732

The predicted product size and location of oligonucleotide binding sites for neuroglobin RT-PCR products are indicated. For neuroglobin RT-PCR products predicted size of product and location of oligonucleotide binding sites are based upon a partial cDNA sequence from turtle brain and the mouse coding sequence (Accession No. NM022414). For actin RT-PCR products predicted size and location of oligonucleotide binding sites are based on the salmon β -actin coding sequence (Accession No. AF309819). Primers for β -actin were homologous to both the salmon and human coding sequences.




Figure 1. Representative gels showing increased neuroglobin transcription relative to actin controls in the *T. scripta* brain. Gels were visualized with ethidium bromide and digitally photographed for analysis. N = normoxia, H = 4 h hypoxia, 1A = 1 h anoxia, 4A = 4 h anoxia, R = 4 h anoxia/4 h reoxygenation.




Figure 2. Changes in neuroglobin mRNA in the hypoxic and anoxic turtle brain, expressed as percent of normoxic control. Each experiment utilized five individuals per group. N = normoxia, H = 4 h hypoxia, 1A = 1 h anoxia, 4A = 4 h anoxia, R = 4 h anoxia/4 h reoxygenation. Columns are numbered 1–5. Column 1 (N) differs significantly from column 2 (H), ($p < 0.01$); column 1 (N) differs significantly from column 4 (4A), ($p < 0.05$); column 4 (4A) differs significantly from column 5 (R), ($p < 0.01$).

mRNA expression was seen with a 4.7-fold increase relative to normoxic control levels.

There are two distinct phases in anoxic responses in the turtle brain: the initial entry phase characterized by K_{ATP} channel activation and downregulation of ATP consuming processes and a second long-term (h/days) maintenance of a

deep hypometabolic state [29]. The purpose of this study was to establish whether neuroglobin mRNA expression was upregulated during the long-term maintenance phase, to compare anoxic mRNA expression levels with those obtained over the same duration (4 h) of hypoxia, and to establish whether any further upregulation occurred upon reoxygenation after anoxia. Previous studies have demonstrated that during 4 h anoxia and with anoxia followed by 4 h reoxygenation there is no alteration in either actin or HIF-1 transcripts confirming that there is no generalized transcriptional response to anoxia or anoxia plus reoxygenation [17].

Discussion

This is the first report of regulated expression of neuroglobin mRNA in response to altered oxygen availability in an anoxia-tolerant organism. The turtle brain affords us the unique opportunity to examine responses to oxygen deprivation in the absence of pathological processes associated with hypoxic responses in mammalian neurons. The upregulation of neuroglobin transcripts during anoxia and during hypoxia may reflect its potential role as a hypoxia sensor, or may relate to an alternative function such as an intracellular oxygen carrier or a component in a detoxification pathway [23, 22, 24].

Any of these functions, however, would increase cell survival under hypoxic conditions. While the upregulation of Ngb by hypoxia and increased neuronal survival *in vitro* and *in vivo* has been reported [25, 27], other investigators, however, found no significant changes in Ngb levels in murine brains after chronic hypoxic (10% O_2) exposure [26]. As earlier work involved anoxic/reperfusion and ischemic animal models [27], however, this difference may simply imply that Ngb is upregulated only under severely hypoxic/anoxic conditions.

Until now the majority of biochemical studies on Ngb have been carried out on mammalian species. However, a comparative approach involving non-mammalian species offers a promising tool for the identification of conserved Ngb features and thus for the further understanding of Ngb function. The advantage of the turtle model is that *in vivo* experiments can be performed on severely

hypoxic and anoxic animals, as they are able to survive and recover from long-term anoxia [for review, see 29]. In this study, 4 h hypoxia (5% O₂) resulted in a more than 3-fold induction of Ngb in the turtle brain, with an even larger increase upon reoxygenation following anoxia. Acute (1 h) and chronic (4 h) anoxia, however, resulted in a much smaller degree of induction, implying that the role or regulation of Ngb in the turtle brain is relatively specific to hypoxia, and that induction is not a rapid process in response to low oxygen levels. Blood gas analysis has demonstrated that blood oxygen levels have essentially reached 0 Torr by the end of the first hour anoxia in vivo [30], however until this time oxygen is present though decreasing. Any rapid, oxygen sensing response of Ngb, then, should have also occurred during the first hour of anoxia in these experiments, however, Ngb mRNA only increased approximately 60% by 1 h anoxia, with a continued increase to 200% of basal by 4 h anoxia despite the complete absence of systemic oxygen.

In hypoxic rat brain, neuroglobin is induced by hypoxia either through HIF-1 signaling or potentially through non-HIF-1 mechanisms. In the rat these changes occur in the context of ongoing pathological neuronal damage. The induction of Ngb by hypoxia in mice, but not by pharmacological agents such as cobalt chloride or deferoxamine [25] suggests the specific regulation of Ngb through hypoxia signaling pathways (HIF-1- α), though alternative regulatory pathways have also been suggested. A comparison of mouse and human Ngb genes, for example, did not show conserved hypoxia responsive elements (HREs) that would permit induction by HIF-1 [31]. While the magnitude of the hypoxic induction in the turtle brain was relatively large the induction in anoxia was more modest. This difference between hypoxic and anoxic responses in turtle brain may suggest either a distinct transcriptional regulatory mechanism or some additional signaling cross talk in the hypoxic state in the turtle brain. Indeed in mammals hypoxic induction of neuroglobin transcripts is induced via a MAP kinase pathway and different MAP kinases are renowned for demonstrating cross talk between each other as well as with other non-MAP kinase pathways [32].

The greater upregulation of Ngb in hypoxia and in anoxia/ reoxygenation, by comparison to anoxia, also implies a functional role in the

presence of oxygen that is not required in anoxia (e.g. ROS detoxification vs. the anoxic regeneration of NAD⁺). As *T. scripta* depends upon metabolic suppression and anaerobic lactate production for anoxic survival, a role for Ngb in the regeneration of NAD⁺ would be likely to be redundant. The absence of a strong upregulation of Ngb under conditions of no oxygen rather than when oxygen is present would also argue against a role for Ngb in the regeneration of NAD⁺.

As previously reported [21] the neuroprotective role of elevated Ngb expression could also be due to the degradation of ROS, though no study has yet verified this possibility. Neurons consume large amounts of oxygen, and are particularly vulnerable to ROS damage due to the abundance of unsaturated fatty acids, which propagate lipid peroxidation, and low ROS defenses [33]. The presence of Ngb in highly active tissues including fish gills [34], the retina [35] and endocrine tissues [36] could indicate roles in either ROS metabolism or oxygen delivery. The greater induction of Ngb by hypoxia over anoxia, and additional increases upon reoxygenation, may indicate a more likely role as an ROS scavenger in the turtle brain than as a means to increase oxygen delivery.

In conclusion, we have observed a major induction of neuroglobin gene expression in hypoxia and upon post-anoxic-reoxygenation in the brain of the anoxia-tolerant turtle. A smaller upregulation of neuroglobin gene expression was found during sustained anoxia. These observations point to a key protective role for neuroglobin in scavenging of reactive oxygen species under conditions of oxygen deprivation and upon reoxygenation.

Acknowledgements

This work was supported by the Florida Atlantic University Foundation, and the American Heart Association (Florida Affiliate). This is the contribution number P200523 from the Florida Center of Excellence in Biomedical and Marine Biotechnology.

References

1. Lutz P.L., Prentice H.M. and Nilsson G.E., *The Brain without Oxygen*, 3rd ed. Kluwers Press, Netherlands, 2003.

2. Liu J., Narasimhan P., Yu F. and Chan P.H., Neuroprotection by hypoxic preconditioning involves oxidative stress-mediated expression of hypoxia-inducible factor and erythropoietin. *Stroke* 36(6), 1264–1269, 2005.
3. Date T., Mochizuki S., Belanger A.J., Yamakawa M., Luo Z., Vincent K.A., Cheng S.H., Gregory R.J. and Jiang C., Expression of constitutively stable hybrid hypoxia-inducible factor-1 α protects cultured rat cardiomyocytes against simulated ischemia-reperfusion injury. *Am. J. Physiol. Cell Physiol.* 288(2), C314–C320, 2005.
4. Alsbury S., Papageorgiou K. and Latchman D.S., Heat shock proteins can protect aged human and rodent cells from different stressful stimuli. *Mech. Ageing Dev.* 125(3), 201–209, 2004.
5. Yamada K. and Inagaki N., Neuroprotection by KATP channels. *J. Mol. Cell Cardiol.* 38(6), 945–949, 2005.
6. Zhao P., Qian H. and Xia Y., GABA and glycine are protective to mature but toxic to immature rat cortical neurons under hypoxia. *Eur. J. Neurosci.* 22(2), 289–300, 2005.
7. Germack R. and Dickenson J.M., Adenosine triggers preconditioning through MEK/ERK1/2 signalling pathway during hypoxia/reoxygenation in neonatal rat cardiomyocytes. *J. Mol. Cell Cardiol.* 39(3), 429–442, 2005.
8. Tomaselli B., Podhraski V., Heftberger V., Bock G. and Baier-Bitterlich G., Purine nucleoside-mediated protection of chemical hypoxia-induced neuronal injuries involves p42/44 MAPK activation. *Neurochem. Int.* 46(7), 513–521, 2005.
9. Jackson D.C., Living without oxygen: lessons from the freshwater turtle. *Comp. Biochem. Physiol. A* 125(3), 299–315, 2000.
10. Milton S.L. and Lutz P.L., Low extracellular dopamine levels are maintained in the anoxic turtle brain. *J. Cereb. Blood Flow Metab.* 18: 803–807, 1998.
11. Milton S.L., Thompson J.W. and Lutz P.L., Mechanisms for maintaining extracellular glutamate levels in the anoxic turtle striatum. *J. Appl. Physiol.* 282: R1317–R1323, 2002.
12. Nilsson G.N. and Lutz P.L., Adenosine release in the anoxic turtle brain: a possible mechanism for anoxic survival. *J. Exp. Biol.* 162: 345–351, 1992.
13. Nilsson G.N. and Lutz P.L., Release of inhibitory neurotransmitters in response to anoxia in turtle brain. *Am. J. Physiol. Reg. Integr. Comp. Physiol.* 261: R32–R37, 1991.
14. Fernandes J.A., Lutz P.L., Tannenbaum A., Todorov A.T., Liebovitch L. and Vertes R., Electroencephalogram activity in the anoxic turtle brain. *Am. J. Physiol.* 273: R 911–R919, 1997.
15. Greenway S.C. and Storey K.B., Mitogen-activated protein kinases and anoxic tolerance in turtles. *J. Exp. Zool.* 287(7), 477–484, 2000.
16. Prentice H.M., Milton S.L., Scheurle D. and Lutz P.L., The upregulation of cognate and inducible heat shock proteins in the anoxic turtle brain. *J. Cereb. Blood Flow Metab.* 24: 826–828, 2004.
17. Prentice H.M., Milton S.L., Scheurle D. and Lutz P.L., Gene transcription of brain voltage-gated potassium channels is reversibly regulated by oxygen supply. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 285(6), R1317–R1321, 2003.
18. Belkin D.A., Anoxia: tolerance in reptiles. *Science* 139: 492–493, 1963.
19. Nilsson G.N., Lutz P.L. and Jackson T.L., Neurotransmitters and anoxic survival in the brain: A comparison of anoxia-tolerant and anoxia-intolerant vertebrates. *Physiol. Zool.* 64(3), 638–652, 1991.
20. Burmester T., Weich B., Reinhardt S. and Hankeln T., A vertebrate globin expressed in the brain. *Nature* 407: 520–523, 2000.
21. Burmester T. and Hankeln T., Neuroglobin: A respiratory protein of the nervous system. *News Physiol. Sci.* 19: 110–113, 2004.
22. Sowa A.W., Duff S.M.G., Guy P.A. and Hill R.D., Altering hemoglobin levels changes energy status in maize cells under hypoxia. *PNAS* 95: 10317–10321, 1998.
23. Herold S., Fago A., Weber R.E., Dewilde S. and Moens L., Reactivity studies of the Fe(III) and Fe(II)NO forms of human neuroglobin reveal a potential role against oxidative stress. *J. Biol. Chem.* 279: 22841–22847, 2004.
24. Kriegl J.M., Bhattacharyya A.J., Nienhaus K., Deng P., Minkow O. and Nienhaus G.U., Ligand binding and protein dynamics in neuroglobin. *PNAS* 99: 7992–7997, 2002.
25. Sun Y., Jin K., Mao X.O., Zhu Y. and Greenberg D.A., Neuroglobin is upregulated by and protects neurons from hypoxic-ischemic injury. *PNAS USA* 98: 15306–15311, 2001.
26. Mammem P.P.A., Shelton J.M., Goetsch S.C., Williams S.C., Richardson J.A., Garry M.G. and Garry D.J., Neuroglobin, a novel member of the globin family, is expressed in focal regions of the brain. *J. Histochem. Cytochem.* 50(12), 1591–1598, 2002.
27. Sun Y., Jin K., Peel A., Mao X.O., Xie L. and Greenberg D.A., Neuroglobin protects the brain from experimental stroke *in vivo*. *PNAS USA* 100: 3497–3500, 2003.
28. Bickler P.E. and Donohoe P.H., Adaptive responses of vertebrate neurons to hypoxia. *J. Exp. Biol.* 205(Pt 23), 3579–3586, 2002.
29. Lutz P.L. and Milton S.L., Negotiating brain anoxia survival in the turtle. *J. Exp. Biol.* 207: 3141–3147, 2004.
30. Milton, S.L., The physiology of hypoxia and anoxia tolerance in three species of turtle: the loggerhead sea turtle (*Caretta caretta*), green sea turtle (*Chelonia mydas*) and freshwater *Trachemys scripta*. Ph.D. Thesis, University of Miami, 1994, pp 196.
31. Wystub S., Laufs T., Schmidt M., Burmester T., Maas U., Saaler-Reinhardt S., Hankeln T. and Reuss S., Localization of neuroglobin protein in the mouse brain. *Neurosci. Lett.* 346: 114–116, 2003.
32. Zhu Y., Sun Y., Kunlin J. and Greenberg D.A., Hemin induces neuroglobin expression in neural cells. *Blood* 100: 2494–2498, 2002.
33. Evans P.H., Free radicals in brain metabolism and pathology. *Br. Med. Bull.* 49: 577–587, 1993.
34. Fuchs C., Heibs V., Kiger L., Haberkamp M., Roesner A., Schmidt M., Hamdane D., Marden M.C., Hankeln T. and Burmester T., Zebrafish reveals different and conserved features of vertebrate neuroglobin gene structure, expression pattern, and ligand binding. *J. Biol. Chem.* 279: 24116–24122, 2004.
35. Schmidt M., Giel A., Laufs T., Hankeln T., Wolfrum U. and Burmester T., How does the eye breathe? Evidence for neuroglobin-mediated oxygen supply of the mammalian retina. *J. Biol. Chem.* 278: 1932–1935, 2003.
36. Reuss S., Saaler-Reinhardt S., Weich B., Wystub S., Reuss M., Burmester T. and Hankeln T., Expression analysis of neuroglobin mRNA in rodent tissues. *Neuroscience* 115: 645–656, 2002.